

# Study guide for exam #2

Chemistry 342, (01:160:342), Spring 2012  
Physical Chemistry of Biochemical Systems

## 1 Key concepts and equations

For the following, you should understand the concepts, know (have memorized) the key equations, understand what all the symbols mean, and be able to explain the equations in words and to appreciate their context. For chapters 5-7, please pay close attention to the *Checklist of key concepts* and the *Checklist of key equations*.

### 1.1 Chapter 5

- mean activity coefficients,  $\gamma_{\pm} = (\gamma_+^p \gamma_-^q)^{1/s}$ ;  $s = p + q$
- ionic strength:  $I = \frac{1}{2} \sum z_i^2 b_i$
- transfer across a membrane:  $\Delta G_m = RT \ln(a_{in}/a_{out}) + zF \Delta \phi$
- Nernst equation:  $E_{cell} = E_{cell}^{\ominus} - (RT/vF) \ln Q$
- dependence of cell potential on pH:  $E = E' - \frac{v_{H^+} RT \ln 10}{v_e F} \times pH$

### 1.2 Chapter 6

- differences between differential and integrated rate laws
- half life for a non-reversible first-order reaction:  $t_{1/2} = (\ln 2)/k_r$
- Arrhenius equation:  $k_r = \exp(-E_a/RT)$

### 1.3 Chapter 7

- relation of equilibrium and rate constants:  $K = k_r/k_r'$
- relaxation times:  $x = x_0 \exp(-t/\tau)$ ;  $\tau^{-1} = k_r + k_r'$
- diffusion controlled reaction:  $k_d = (8RT/3\eta)$  [you don't need to memorize this one!]

## 2 Sample exam questions

1. Explain (define in words) each of the symbols in the above expressions; explain their limitations (what they assume)
2. Does  $\text{FADH}_2$  have a tendency to reduce coenzyme Q at pH 7? (Use values in Table 5.2).
3. What is the equilibrium constant for the reaction in example 2 at pH 7? at pH 4?
4. Cytochrome c oxidase accepts electrons from reduced cytochrome c ( $\text{Fe}^{+2}\text{cyt c}$ ) and transmits them to molecular oxygen, forming water. Write a balanced reaction for this process, and estimate  $E_{cell}^{\oplus}$ . Does lowering the pH promote or inhibit the reaction? (Use data in Table 5.2)
5. The anesthetic phenobarbital metabolizes following first-order kinetics, with a half-life of 4.5 hours. For an initial dose of 450 mg, how much remains after 2 hours?
6. What are the units of  $k_r$  for zeroth, first, second and third-order reactions?
7. A protein dimerizes  $2A \leftrightarrow A_2$  with a forward rate constant (second order) of  $k_r$  and a reverse rate constant (first order) of  $k'_r$ . Show that the relaxation time is  $\tau^{-1} = k'_r + 4k_r[A_{eq}]$ .